

Genetic characterization of parthenogenic *Fasciola* sp. in Japan on the basis of the sequences of ribosomal and mitochondrial DNA

T. ITAGAKI^{1*}, M. KIKAWA¹, K. SAKAGUCHI¹, J. SHIMO¹, K. TERASAKI², T. SHIBAHARA³ and K. FUKUDA⁴

¹Laboratory of Veterinary Parasitology, Faculty of Agriculture, Iwate University, Ueda 3-18-8, Morioka 020-8550, Japan

²St Mary's Junior College, Kurume 830-8558, Japan

³Division of Laboratory Animal Science, Research Center for Bioscience and Technology, Tottori University, Yonago 683-8503, Japan

⁴Center for Laboratory Animal Science, National Defense Medical College, Tokorozawa 359-8513, Japan

(Received 25 March 2005; revised 9 May 2005; accepted 9 May 2005)

SUMMARY

Accurate identification of aspermic *Fasciola* forms in Japan remains difficult because of their morphological variations. In order to characterize the forms genetically, nucleotide sequences of ribosomal internal transcribed spacer (ITS1 and ITS2) and mitochondrial cytochrome c oxidase I (COI) and NADH dehydrogenase I (NDI) genes in 34 liver flukes from 16 prefectures in Japan were analysed. Two major forms represented by Fsp 1 and Fsp 2 had sequences identical to or closely resembling those of *F. hepatica* and *F. gigantica*, respectively, in all the 4 DNA markers and were mainly distributed in northern and eastern-western parts of Japan, respectively. Fsp 1 and Fsp 2 would have been introduced into Japan with infected cattle of 2 distinct lineages via the Korean Peninsula and spread through limited parts of Japan (northern and eastern-western parts) together with the movement of each cattle lineage. The Japanese form (Fsp 1/2), which showed heterozygosity in ribosomal DNA and Fsp 2 haplotype in mitochondrial DNA, may have originated in interspecific cross hybridization between paternal *F. hepatica* and maternal *F. gigantica*.

Key words: *Fasciola* sp., Japan, parthenogenesis, haplotype, ITS1, ITS2, COI, NDI, genetic characterization.

INTRODUCTION

The two species commonly recognized as the causative agents of fascioliasis in domestic animals and humans are *Fasciola hepatica* and *F. gigantica*. *F. hepatica* mainly occurs in Europe, Americas and Oceania, while *F. gigantica* is distributed in Africa and Asia (Torgerson and Claxton, 1999). Although the two species are classified by morphological characters, especially body length and width, it is difficult accurately to discriminate between the two species because of many variations in size depending on such factors as the age of the flukes, the species of host and the technical difficulty in fixation of the flukes (Kendall, 1965). The *Fasciola* species in Japan had been believed to be *F. hepatica* until Watanabe and Iwata (1954) raised doubt concerning the species. Their report motivated us to carry out research to identify the Japanese species of *Fasciola*. As a result, it has become clear that Japanese *Fasciola* species include worms morphologically resembling *F. hepatica*, *F. gigantica* and their intermediate forms

and that they have ecological characters resembling *F. hepatica* more than *F. gigantica* (Watanabe and Iwata, 1954; Itagaki and Akane, 1959; Oshima, Akahane and Shimazu, 1968; Akahane, Harada and Oshima, 1970). In addition, *Fasciola* species in Japan has been found to have cytogenetical characteristics differing from those of *F. hepatica* and *F. gigantica*. The Japanese forms include worms with 20 or 30 chromosomes corresponding to diploid and triploid sets and with 20/30 mosaic ploidy, and they are meiotically dysfunctional and aspermic (Moriyama, Tsuji and Seto, 1979; Sakaguchi, 1980), while *F. hepatica* and *F. gigantica* are spermic diploid and meiotically functional (Sanderson, 1953; Reddy and Subramanyam, 1973). Aspermic diploids and triploids in Japan had univalent chromosomes in primary oocytes, suggesting the potential of parthenogenesis (Terasaki *et al.* 2000). From these findings, it is still difficult accurately to identify Japanese *Fasciola* species.

Molecular genetic approaches based on DNA analysis have been employed to solve the problems of identification and genetic characterization of morphologically similar parasites, including *Fasciola* species (Blair and McManus, 1989; Adlard *et al.* 1993; McManus and Bowles, 1996). There have been studies on genetic characterization of *Fasciola*

* Corresponding author: Laboratory of Veterinary Parasitology, Faculty of Agriculture, Iwate University, Ueda 3-18-8, Morioka 020-8550, Japan. Tel and Fax: +81 19 621 6219. E-mail: itagaki@iwate-u.ac.jp

species in Japan based on ribosomal and mitochondrial DNA (Hashimoto *et al.* 1997; Itagaki *et al.* 1998; Itagaki and Tsutsumi, 1998). Hashimoto *et al.* (1997) and Itagaki *et al.* (1998) proved that a Japanese form was composed of flukes genetically identical to *F. gigantica*, whereas Itagaki and Tsutsumi (1998) showed the occurrence of those resembling *F. hepatica* and *F. gigantica*. These differences probably result from the small numbers of samples analysed and the lack of information on divergence of ribosomal DNA and that of mitochondrial DNA. In order accurately to characterize the Japanese *Fasciola* forms, we analysed the nucleotide sequences of ribosomal ITS1 and ITS2 and mitochondrial NDI and COI genes using 34 *Fasciola* samples collected in 16 prefectures of Japan.

MATERIALS AND METHODS

Fasciola individuals and chromosome observation

Thirty-four adults of *Fasciola* sp. were obtained from the bile ducts of infected cattle and sika deer (*Cervus nippon centralis* and *C. n. ezoensis*) in 16 different prefectures of Japan and kept at -80°C or in 70% ethanol until DNA extraction. In some of the flukes, a small piece of the testis was cut off and fixed in freshly prepared ethanol-acetic acid (3:1) before freezing or fixation with 70% ethanol, and chromosomes were observed by using a squashing method (Snow, 1963). *F. hepatica* and *F. gigantica* that had been precisely identified by morphological features, including ratio of body length and width and the presence of sperm within the vesicula seminalis, were used for comparing degrees of DNA divergence. The *F. hepatica* used was from Uruguay (2 specimens), Australia (2) and Northern Ireland (2), and the *F. gigantica* used were from Zambia (6), Thailand (2) and Indonesia (3).

DNA extraction and amplification

Total DNA was extracted from individual flukes by using an E.Z.N.A. mollusk DNA kit (Omega Bio-tek, Doraville, USA). DNA fragments of each target region were amplified by PCR using 1.25 units of *Taq* polymerase (Promega, Madison, USA), 0.4 mM each of dATP, dTTP, dCTP and dGTP, 2 mM MgCl_2 , each primer set (50 pmol/25 μl reaction mixture), and PCR buffer. The primer sets used to amplify the fragments were Ita 10 (5'-AAGG-ATGTTGCTTTGTCGTGG-3') and Ita 2 (5'-GGAGTACGGTTACATTCACA-3') for NDI, Ita 8 (5'-ACGTTGGATCATAAGCGTGT-3') and Ita 9 (5'-CCTCATCCAACATAACCTCT-3') for COI, ITS1-F (5'-TTGCGCTGATTACGT-CCCTG-3') and ITS1-R (5'-TTGGCTGCGC-TCTTCATCGAC-3') for ITS1, and ITS2-F (5'-TGTGTCGATGAAGAGCGCAG-3') and

ITS2-R (5'-TGGTTAGTTTCTTTTCCTCC-GC-3') for ITS2. Reaction cycles consisted of an initial denaturing step at 94°C for 90 sec, followed by 30 cycles at 94°C for 90 sec, 55°C for 90 sec and 72°C for 120 sec with a final extension at 72°C for 10 min using GeneAmp PCR Systems 2400 and 2700 (Applied Biosystems, Tokyo, Japan). PCR products were precipitated with ethanol/sodium acetate and dissolved in MilliQ water.

Sequencing analysis

PCR fragments were directly sequenced using ABI Prism Big Dye terminator v. 1.0 and v. 3.0 ready reaction cycle sequencing kits (Applied Biosystems) with the use of an additional primer, Ita 4 (5'-CTATAACCAGTCATACT-3'), for ND1 together with the same primers as those used in PCR. At least 2 fragments amplified for individual flukes were sequenced for each target region in both directions using forward and reverse primers. The sequencing reactions were run on a PE Applied Biosystems 310 or 3100 automated sequencer. The sequence data were aligned by Clustal X program v. 1.53 b (Thompson *et al.* 1997). Phylogenetic analyses were conducted by neighbour-joining (NJ) and maximum parsimony (MP) using PAUP 4.0b10 (Swofford, 2001) with a lung fluke, *Paragonimus westermani* (Accession no. AF219379) designed as an outgroup. All characters were run unordered and equally weighted. Alignment gaps were treated as missing data. A heuristic search with tree-bisection-reconnection (TBR)-branch swapping was used in MP analysis to infer the shortest trees. The length, consistency index excluding uninformative characters (C.I.) and retention index (R.I.) of the most parsimonious trees were recorded. A bootstrap analysis (using 1000 replicates) was conducted using heuristic searches and TBR branch swapping with the MulTrees option in order to determine the relative support for clades of the consensus tree.

RESULTS

Ploidy

Chromosomes were observed in 12 of the 34 Japanese flukes. The number of chromosomes was counted in well-spread primary spermatocytes. Eleven flukes were found to have 30 univalent chromosomes ($3n=30$, triploid), and one had 20 univalent chromosomes ($2n=20$, diploid) (Table 1).

Haplotype from ITS1 and ITS2 sequences

The sequences of the ITS1 region in *Fasciola* specimens (Accession numbers AB207139-AB207147) consisted of 600 bps including complete ITS1,

Table 1. Haplotype and ploidy of parthenogenic *Fasciola* sp. in Japan

| Specimen code# | Locality | Ploidy | Haplotype | | | |
|----------------|-----------|--------|-----------|---------|-------|-------|
| | | | ITS1 | ITS2 | NDI | COI |
| 1 | Hokkaido | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 2 | Hokkaido | ND | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 3 | Aomori | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 4 | Aomori | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 5 | Iwate | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 6 | Iwate | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 7 | Iwate | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 8 | Tochigi | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 9 | Tochigi | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 10 | Saitama | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 11 | Saitama | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 12 | Nagano | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 13 | Nagano | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 14 | Nagano | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 15 | Nara | 3n | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 16 | Hyogo | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 17 | Hyogo | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 18 | Hiroshima | 2n | Fsp 1/2 | Fsp 1/2 | Fsp 2 | Fsp 2 |
| 19 | Hiroshima | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 20 | Hiroshima | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 21 | Tottori | ND | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 22 | Tottori | ND | Fsp 1 | Fsp 1 | Fsp 1 | Fsp 1 |
| 23 | Tokushima | 3n | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 24 | Tokushima | 3n | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 25 | Kochi | ND | Fsp 2 | Fsp 2a | Fsp 2 | Fsp 2 |
| 26 | Kochi | ND | Fsp 2 | Fsp 2a | Fsp 2 | Fsp 2 |
| 27 | Saga | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 28 | Saga | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 29 | Nagasaki | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 30 | Nagasaki | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 31 | Kumamoto | ND | Fsp 1 | Fsp 1 | Fsp 2 | Fsp 2 |
| 32 | Kumamoto | ND | Fsp 2 | Fsp 2 | Fsp 2 | Fsp 2 |
| 33 | Kagoshima | ND | Fsp 1/2 | Fsp 1/2 | Fsp 2 | Fsp 2 |
| 34 | Kagoshima | ND | Fsp 1/2 | Fsp 1/2 | Fsp 2 | Fsp 2 |

'2n' and '3n' mean diploid and triploid, respectively. ND, no data.

partial 18S and 5·8S rDNA and had 6 variable nucleotide positions of which 1 was located in 18S and 5 were located in ITS1. The sequences of *F. hepatica* and *F. gigantica* differed at the 6 positions. Japanese flukes showed 3 different sequences (haplotypes) represented by Fsp 1, Fsp 2 and Fsp 1/2 which had nucleotides identical to *F. hepatica* and *F. gigantica* and those overlapped between the two species, respectively (Table 2). The 505-bp or 506-bp sequence of the ITS2 region included complete ITS2, partial 5·8S and 28S, and 8 nucleotide positions (including 1 gap) within the sequence were variable among individuals of *Fasciola* spp. (Accession nos. AB207148–AB207153). Japanese flukes were divided into 4 haplotypes (Fsp 1, Fsp 2, Fsp 2a, Fsp 1/2) based on the nucleotides at the 7 sites. Fsp 1 and Fsp 2 were identical or very similar to sequences of *F. hepatica* and *F. gigantica*, respectively. Fsp 2a differed from Fsp 2 in only 1 nucleotide site (331). Fsp 1/2 had nucleotides overlapped between Fsp 1 and Fsp 2 at 5 sites (307, 318,

331, 370, 376) and the same nucleotides as Fsp 1 at 2 sites (424, 434). The number of Japanese flukes that showed the haplotypes of Fsp 1, Fsp 2 including Fsp 2a and Fsp 1/2 in both ITS1 and ITS2 regions were 13, 18 and 3, respectively. All of the flukes obtained from Hokkaido, Aomori, Iwate and Tochigi Prefectures in the northern part of Japan showed Fsp 1 in both regions, whereas most of the flukes from the western (Kagoshima) to eastern (Saitama) parts of Japan showed Fsp 2 (Fig. 1). Eleven triploid flukes showed either Fsp 1 or Fsp 2, and 1 diploid showed Fsp 1/2 (Table 1).

Haplotype from COI and NDI sequences

Partial COI sequence (438 bp) and NDI sequence (535 bp) were determined for 51 *Fasciola* individuals used and included 62 and 68 variable sites, respectively (Accession nos. AB207170–AB207183, AB207103, AB207154–AB207169). Although intra-specific variability in COI and NDI sequences was

Table 2. Comparison of the nucleotides at variable sites in the ITS1 and ITS2 regions of *Fasciola* spp.

| Species | Locality and haplotype | Sites of ITS1 region | | | | | | Sites of ITS2 region | | | | | | No. of worms | | |
|---------------------|------------------------|----------------------|-----|-----|-----|-----|-----|----------------------|-----|-----|-----|-----|-----|--------------|-----|-----|
| | | 48 | 175 | 265 | 359 | 437 | 457 | 307 | 318 | 331 | 370 | 376 | 384 | | 424 | 434 |
| <i>F. hepatica</i> | Uruguay | T | C | A | C | T | C | T | T | T | C | C | T | T | G | 2 |
| | Australia | T | C | A | C | T | C | T | T | T | C | C | C | T | G | 2 |
| | N. Ireland | T | C | A | C | T | C | T | T | T | C | C | C | T | G | 2 |
| <i>F. gigantica</i> | Indonesia | C | T | T | T | A | T | C | T | C | T | T | C | — | A | 2 |
| | Thailand | C | T | T | T | A | T | C | T | C | T | T | C | — | A | 2 |
| | Zambia | C | T | T | T | A | T | T | T | C | T | T | C | — | A | 6 |
| <i>Fasciola</i> sp. | Japan; Fsp 1 | T | C | A | C | T | C | T | T | T | C | C | C | T | G | 13 |
| | Japan; Fsp 2 | C | T | T | T | A | T | C | C | C | T | T | C | — | A | 16 |
| | Japan; Fsp 2a | C | T | T | T | A | T | C | C | T | T | T | C | — | A | 2 |
| | Japan; Fsp 1/2 | T/C | C/T | A/T | C/T | T/A | C/T | T/C | T/C | T/C | C/T | C/T | C | T | G | 3 |




Fig. 1. Distribution of the haplotypes of *Fasciola* sp. in Japan. White, black and striped circles represent the haplotypes, Fsp 1, Fsp 2 and Fsp 1/2, respectively.

found in *F. hepatica* and *F. gigantica*, the degree of variability was higher in *F. gigantica* than in *F. hepatica*. Japanese specimens had either of 2 distinct sequences (haplotypes: Fsp 1, Fsp 2) in both COI and NDI. Fsp 1 and Fsp 2 differed from each other at 41 nucleotide positions within COI and at 45 sites within NDI. Fsp 1 was very similar to the sequence of *F. hepatica* in both COI and NDI, though no complete consistency was confirmed between the sequences. In contrast, Fsp 2 was similar to the sequence of *F. gigantica*, being more similar to the sequence of *F. gigantica* obtained from Indonesia and Thailand than from Zambia. Twelve of 34 flukes showed Fsp 1, and the other 22 showed Fsp 2. MP analysis resolved most parsimonious trees for COI (length 356; CI=0.9270; RI=0.9323) and NDI (length 522; CI=0.8966; RI=0.8893)

(Fig. 2). Tree topologies from MP and NJ (not shown) analyses were consistent with each other in both sequence sets of COI and NDI. The individuals of *F. hepatica* and Fsp 1 belonged to the same clade, which clearly differed from the clade of *F. gigantica* and Fsp 2.

DISCUSSION

The haplotypes based on the sequences of ITS1, ITS2, COI and NDI showed that 2 major and 3 minor distinct forms of *Fasciola* occurred in Japan. Similarly, 3 different forms identified using isozyme patterns have been found in Japanese triploid *Fasciola* (Agatsuma *et al.* 1994), suggesting at least 3 origins of triploidy. Recently, Terasaki *et al.* (2000) have suggested that the most likely origin of triploid




Fig. 2. Maximum parsimony trees of *Fasciola* spp. inferred from the nucleotide sequences of partial NDI (A) and COI (B). Fsp 1 and Fsp 2 show the haplotypes of *Fasciola* sp. in Japan. *F. h.* and *F. g.* show *F. hepatica* and *F. gigantica*, respectively. '#' and parentheses represent code no. and locality of individual, respectively.

forms was the fusion of a sperm of a bisexual diploid (*F. hepatica* or *F. gigantica*) with an unreduced egg of a parthenogenetic diploid. According to this hypothesis, the triploid specimens (e.g., #1) that showed the genotype of Fsp 1 (*F. hepatica* type) in both ribosomal and mitochondrial DNA would be derived from hybridization between parental *F. hepatica* and parthenogenetic diploid (*F. hepatica* type). In fact, parthenogenetic diploid forms closely resembling *F. hepatica* in sequences of rDNA and mtDNA have been found in Korea (Itagaki *et al.*, unpublished data). Similarly, the triploid individuals (e.g., #23) with the genotype of Fsp 2 in both markers seem to be crossing progenies between *F. gigantica* and parthenogenetic diploid (*F. gigantica* type). On the other hand, the parthenogenetic diploid appears to be directly derived from bisexual *F. hepatica* or *F. gigantica* populations (Terasaki *et al.* 2000). Furthermore, in most animal taxa, the acquisition of clonal reproduction may be a consequence of hybridization between sexual species (Dawley, 1989; Wilson and Hebert, 1992; Vrijenhoek, 1994). Most clonal lineages therefore contain a mixture of the nuclear genomes of the two or more sexual species (Avisé, Quattro and Vrijenhoek, 1992). Thus, the aspermic diploid (#18)

that showed a heterozygous genotype (Fsp 1/2) in rDNA and Fsp 2 in mtDNA may have originated in interspecific cross-hybridization between paternal *F. hepatica* and maternal *F. gigantica*. This natural hybridization between the two bisexual *Fasciola* species has been also suggested by results of previous analyses (Itagaki and Tsutsumi, 1998; Agatsuma *et al.* 2000).

Although aspermic (parthenogenetic) *Fasciola* is widely distributed in East and Southeast Asia (Terasaki, Akahane and Habe, 1982), the origin still remains unknown. However, parthenogenetic *Fasciola* would not have originated in Japan as well as in Korea, since no bisexual *Fasciola* has been found in those countries. We have recently found that triploid individuals 100% identical to Fsp 2 in the sequences of not only ITS1 but also mitochondrial COI and NDI with rapid base substitution were present in Vietnam, which is far from Japan (Itagaki *et al.*, unpublished data). The origin of the triploid forms, however, is also not Vietnam, since no parthenogenetic diploid has been found in Vietnam. These findings strongly suggest that the triploid forms with the genotype of Fsp 2 originated in a country other than Japan, Korea and Vietnam and rapidly spread among these countries with infected

animals (probably domestic cattle). Molecular phylogeny using *Fasciola* individuals from other Asian countries is needed to elucidate the origin and route of spread throughout Asia of parthenogenic *Fasciola* forms.

Japanese native cattle were thought to have been introduced into Japan via the Korean Peninsula in about the second century AD (Mukai, 1989). In fact, Japanese and Korean native cattle consist of haplotypes very similar to each other in mtDNA sequences. Interestingly, only 2 of the haplotypes are major in Japan (Mannen *et al.* 2004). Furthermore, *Fasciola* forms that are completely identical to Fsp 1 and Fsp 2 in sequences of both rDNA and mtDNA occur in Korea (Itagaki *et al.*, unpublished data). These findings strongly suggest that Fsp 1 and Fsp 2 invaded Japan with infected cattle and spread through limited parts of Japan together with the movement of the 2 haplotypes of cattle.

This study was supported in part by a Grant-in-Aid for Scientific Research (C) (no. 14560243) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

REFERENCES

- Adlard, R. D., Barker, S. C., Blair, D. and Cribb, T. H. (1993). Comparison of the second internal transcribed spacer (ribosomal DNA) from populations and species of Fasciolidae (Digenea). *International Journal for Parasitology* **23**, 423–425.
- Agatsuma, T., Terasaki, K., Yang, L. and Blair, D. (1994). Genetic variation in the triploids of Japanese *Fasciola* species, and relationships with other species in the genus. *Journal of Helminthology* **68**, 181–186.
- Agatsuma, T., Arakawa, Y., Iwagami, M., Honzako, Y., Cahyaningsih, U., Kang, S-Y. and Hong, S-J. (2000). Molecular evidence of natural hybridization between *Fasciola hepatica* and *F. gigantica*. *Parasitology International* **49**, 231–238.
- Akahane, H., Harada, Y. and Oshima, T. (1970). Patterns of variation of the common liver fluke (*Fasciola* sp.) in Japan. III. Comparative studies on the external form, size of egg and number of eggs in the uterus of fluke in cattle, goat and rabbit. *Japanese Journal of Parasitology* **19**, 619–627 (in Japanese with English summary).
- Avise, J. C., Quattro, J. M. and Vrijenhoek, R. C. (1992). Molecular clones within organismal clones: mitochondrial DNA phylogenies and the evolutionary histories of unisexual vertebrates. *Journal of Evolutionary Biology* **26**, 225–246.
- Blair, D. and McManus, D. P. (1989). Restriction enzyme mapping of ribosomal DNA can distinguish between fasciolid (liver fluke) species. *Molecular and Biochemical Parasitology* **36**, 201–208.
- Dawley, R. M. (1989). An introduction to unisexual vertebrates. In *Evolution and Ecology of Unisexual Vertebrates* (ed. Dawley, R. M. and Bogart, J. P.), pp. 1–18. New York State Museum, Albany, New York.
- Hashimoto, K., Watanobe, T., Liu, C. X., Init, I., Blair, D., Ohnishi, T. and Agatsuma, T. (1997). Mitochondrial DNA and nuclear DNA indicate that the Japanese *Fasciola* species is *F. gigantica*. *Parasitology Research* **83**, 220–225.
- Itagaki, H. and Akane, S. (1959). Morphological study on the Japanese liver fluke, compared with the African specimens. *The Bulletin of Azabu Veterinary College* **6**, 115–123.
- Itagaki, T. and Tsutsumi, K. (1998). Triploid form of *Fasciola* in Japan: genetic relationships between *Fasciola hepatica* and *Fasciola gigantica* determined by ITS-2 sequence of nuclear rDNA. *International Journal for Parasitology* **28**, 777–781.
- Itagaki, T., Tsutsumi, K., Ito, K. and Tsutsumi, Y. (1998). Taxonomic status of the Japanese triploid forms of *Fasciola*: comparison of mitochondrial ND1 and COI sequences with *F. hepatica* and *F. gigantica*. *Journal of Parasitology* **84**, 445–448.
- Kendall, S. B. (1965). Relationships between the species of *Fasciola* and their molluscan hosts. *Advances in Parasitology* **3**, 59–95.
- Mannen, H., Kohno, M., Nagata, Y., Tsuji, S., Bradley, D. G., Yeo, J. S., Nyamsamba, D., Zagdsuren, Y., Yokohama, M., Nomura, K. and Amano, T. (2004). Independent mitochondrial origin and historical genetic differentiation in North Eastern Asian cattle. *Molecular Phylogenetics and Evolution* **32**, 539–544.
- McManus, D. P. and Bowles, J. (1996). Molecular genetic approaches to parasite identification: their value in diagnostic parasitology and systematics. *International Journal for Parasitology* **26**, 687–704.
- Moriyama, N., Tsuji, M. and Seto, T. (1979). Three karyotypes and their phenotypes of Japanese liver flukes (*Fasciola* sp.). *Japanese Journal of Parasitology* **28**, 23–33 (in Japanese with English summary).
- Mukai, F., Tsuji, S., Fukazawa, K., Ohtagaki, S. and Nambu, Y. (1989). History and population structure of a closed strain of Japanese Black cattle. *Journal of Animal Breeding and Genetics* **106**, 254–264.
- Oshima, T., Akahane, H. and Shimazu, T. (1968). Patterns of the variation of the common liver fluke (*Fasciola* sp.) in Japan. I. Variations in the sizes and shapes of the worms and eggs. *Japanese Journal of Parasitology* **17**, 97–105 (in Japanese with English summary).
- Reddy, P. V. and Subramanyam, S. (1973). Chromosome studies in the liver fluke, *Fasciola gigantica* Cobbold, 1856, from Andhra Pradesh. *Current Science* **42**, 288–291.
- Sakaguchi, Y. (1980). Karyotype and gametogenesis of the common liver fluke, *Fasciola* sp., in Japan. *Japanese Journal of Parasitology* **29**, 507–513.
- Sanderson, A. R. (1953). Maturation and probable gynogenesis in the liver fluke, *Fasciola hepatica* L. *Nature, London* **172**, 110–112.
- Snow, R. (1963). Alcoholic hydrochloric acid-carmines as a stain for chromosomes in squash preparations. *Stain Technology* **38**, 9–13.
- Swofford, D. L. (2001). PAUP*. Phylogenetic analysis using parsimony and other methods ver. 4.0beta. Sinauer Associates, Sunderland, Massachusetts.

- Terasaki, K., Akahane, H. and Habe, S.** (1982). The geographical distribution of common liver flukes (the Genus *Fasciola*) with normal and abnormal spermatogenesis. *Japanese Journal of Parasitology* **44**, 223–231.
- Terasaki, K., Noda, Y., Shibahara, T. and Itagaki, T.** (2000). Morphological comparisons and hypotheses on the origin of polyploids in parthenogenetic *Fasciola* sp. *Journal of Parasitology* **86**, 724–729.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. and Higgins, D. G.** (1997). The Clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acid Research* **24**, 4876–4882.
- Torgerson, P. and Claxton, J.** (1999). Epidemiology and control. In *Fasciolosis* (ed. Dalton, J. P.), pp. 113–149. CABI Publishing, UK.
- Vrijenhoek, R. C.** (1994). Unisexual fish: model systems for studying ecology and evolution. *Annual Review of Ecology and Systematics* **25**, 71–96.
- Watanabe, S. and Iwata, S.** (1954). *Fasciola* species in Japan. *Journal of Japan Veterinary Medical Association* **7**, 124–126 (in Japanese).
- Wilson, C. C. and Hebert, P. D. N.** (1992). The maintenance of taxon diversity in an asexual assemblage: an experimental analysis. *Ecology* **73**, 1462–1472.